

On the limits of Brans-Dicke spacetimes: a coordinate-free approach

F. M. Paiva^{*} and C. Romero[†]

^{*} Centro Brasileiro de Pesquisas Físicas,
R. Dr. Xavier Sigaud 150,
22290-180 Rio de Janeiro – RJ, Brazil

[†] CP 5008 Departamento de Física
Universidade Federal da Paraíba,
58059-900 João Pessoa – PB, Brazil

Abstract

We investigate the limit of Brans-Dicke spacetimes as $\omega \rightarrow \infty$ applying a coordinate-free technique. We obtain the limits of some known exact solutions. It is shown that these limits may not correspond to similar solutions in the general relativity theory.

PACS numbers: 04.20.Jb, 04.20.Cv, 04.50.+h

1 Introduction

In a recent paper [21] it has been showed that as the scalar field coupling constant $\omega \rightarrow \infty$ some solutions of the Brans-Dicke equations for a specified matter configuration (described by the energy-momentum tensor $T_{\alpha\beta}$) do not

^{*}INTERNET: FMIPAIVA@CBPFSU1.CAT.CBPF.BR; BITNET: FMIPAIVA@BRLNCC

[†]BITNET: CENDIF48@BRUFPB

go over to a general relativity solution for the *same* $T_{\alpha\beta}$. This fact raises some discussion about the idea found in the literature (see, for example, [24]) that the Brans-Dicke theory should reduce to general relativity as $\omega \rightarrow \infty$.

Limits of spacetimes as some parameter varies were studied by Geroch [5], who called attention to the fact that the limit may depend on the coordinate system used to perform the calculations. In particular he showed that as the mass tends to infinity the Schwarzschild solution may have as its limits the Minkowski space or a Kasner spacetime. More recently Paiva et al. [16] investigated this subject using a coordinate-free method. Although this new approach was developed in the framework of general relativity it can be also applied to investigate the limits of Brans-Dicke spacetimes.

In this paper we consider three exact solutions of Brans-Dicke field equations. Using the coordinate-free technique we find the limits of these solutions as $\omega \rightarrow \infty$. It turns out that these limit spacetimes may be regarded as solutions of the Einstein general relativity field equations for some specific matter tensors. Then we compare these matter tensors with those of the original Brans-Dicke solutions.

In section 2 we describe briefly the coordinate-free approach to the problem of determining limits of spacetimes in general relativity and show how it can be used to study Brans-Dicke spacetimes.

In section 3 we obtain the limits of the Nariai [17, 18, 12] and the O'Hanlon-Tupper [15] solutions. These are Friedmann-Robertson-Walker (FRW) cosmological models with euclidian spatial section in the Brans-Dicke theory of gravity.

In section 4 we find the limits of a static FRW model with closed spatial section, which is a vacuum solution of Brans-Dicke equations with cosmological constant, found by Romero and Barros [22].

Finally, in the last section, we analyse the matter contents of each limit spacetime obtained as solution of the general relativity theory, drawing conclusions about the limits of Brans-Dicke spacetimes as $\omega \rightarrow \infty$.

2 The coordinate-free approach in Brans-Dicke's theory

The Brans-Dicke [2] theory of gravity assumes, as general relativity does, a lorentzian riemannian manifold structure for the spacetime. The metric is determined through the Brans-Dicke equations

$$R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R = \frac{8\pi}{\phi}T_{\alpha\beta} + \lambda g_{\alpha\beta} + \frac{\omega}{\phi^2} \left(\phi_{,\alpha}\phi_{,\beta} - \frac{1}{2}g_{\alpha\beta}\phi_{,\sigma}\phi^{,\sigma} \right) + \frac{1}{\phi} (\phi_{,\alpha;\beta} - g_{\alpha\beta}\square\phi) \quad (2.1)$$

and

$$\square\phi \stackrel{\text{def}}{=} \phi_{;\alpha}{}^{\alpha} = \frac{(8\pi T + 2\lambda\phi)}{2\omega + 3}, \quad (2.2)$$

where λ is the cosmological constant, ϕ is the scalar field (which plays the role of the inverse of a varying gravitational “constant”), ω is a parameter to be determined “a posteriori”, $T_{\alpha\beta}$ is the matter tensor and T is its trace. To preserve the equivalence principle it is assumed that ϕ has no direct influence on the motion of test particles, which does take place along geodesic lines. Formally, eqs. (2.1) may be regarded as identical to the field equations of general relativity if we define an “effective” energy-momentum tensor $\tau_{\alpha\beta}$ by

$$\frac{8\pi}{G}\tau_{\alpha\beta} = \frac{8\pi}{\phi}T_{\alpha\beta} + \frac{\omega}{\phi^2} \left(\phi_{,\alpha}\phi_{,\beta} - \frac{1}{2}g_{\alpha\beta}\phi_{,\sigma}\phi^{,\sigma} \right) + \frac{1}{\phi} (\phi_{,\alpha;\beta} - g_{\alpha\beta}\square\phi) \quad (2.3)$$

subjected to equation (2.2). Therefore the coordinate-free formalism used in [16] can be used here. In this section we briefly review the main features of this formalism.

With the aim of solving the equivalence problem of metrics, Cartan [3] showed that a spacetime is locally characterized by the components of the Riemann tensor and its covariant derivatives expressed in a constant frame (the *Cartan scalars*), up to (possibly) the 10th order of derivation. Karlhede [8] introduced an algorithm used in practice to calculate the Cartan scalars, which is implemented in the computer algebra system SHEEP [4], and constitute a suite of computer algebra programs called CLASSI [1]. A database of spacetimes together with their Cartan scalars has been produced using CLASSI [14]. To describe this algorithm we introduce the following notation:

R_n : the set of components of the covariant derivatives of the Riemann tensor from the 0th to the n^{th} order — the *Cartan scalars*;

C_n : the algebraically independent Cartan scalars among R_n ;

G_n : the isometry group of R_n ;

H_n : the isotropy group of R_n ;

t_n : the number of independent functions of the coordinates in R_n ;

d_n : the dimension of the orbit of G_n , given by $d_n = 4 - t_n$. Note that $\dim(G_n) = d_n + \dim(H_n)$.

The 0th step of the algorithm is the following. Choose a tetrad basis in which the metric is constant. Calculate R_0 and, using Lorentz transformations, find a basis where it takes a well-defined standard form [8]. Then find H_0 and d_0 . H_0 is found with the help of the Petrov and Segre classifications, while d_0 is the dimension of the spacetime minus t_0 . The next n steps are similar to each other, as described below.

Calculate R_n and, using H_{n-1} , find a basis where R_n takes a well-defined standard form [8]. Then find H_n and d_n . Now, H_n is easily found as it must be a subgroup of H_{n-1} , which is usually of low dimension. If $H_n \subset H_{n-1}$ (strictly, i.e. $H_n \neq H_{n-1}$) or $d_n < d_{n-1}$, the algorithm continues for another step. If $H_n = H_{n-1}$ and $d_n = d_{n-1}$, then R_n can be expressed as functions of R_{n-1} , and so the algorithm stops. In most cases the algorithm stops at the 2nd or 3rd covariant derivative — the 4th is the highest derivative necessary until now [10].

Not all components of the Riemann tensor and its covariant derivatives are algebraically independent; working with spinors MacCallum and Åman [13] found a minimal set of algebraically independent quantities which are actually calculated in the algorithm. Their components will be called *algebraically independent Cartan scalars* and denoted by C_n , according to the order of derivation. For conformally flat solutions, as is the case of the present paper, C_2 is:

$$0^{\text{th}} \text{ derivative : } \quad \Phi_{AB'}, \quad A, B = (0, 1, 2); \quad (2.4)$$

$$\Lambda; \quad (2.5)$$

$$1^{\text{st}} \text{ derivative : } \quad \nabla \Phi_{AB'}, \quad A, B = (0, \dots, 3); \quad (2.6)$$

$$\nabla\Lambda_{AB'}, \quad A, B = (0, 1); \quad (2.7)$$

$$2^{\text{nd}} \text{ derivative : } \nabla^2\Phi_{AB'}, \quad A, B = (0, \dots, 4); \quad (2.8)$$

$$\square\Phi_{AB'}, \quad A, B = (0, 1, 2); \quad (2.9)$$

$$\square\Lambda \quad (2.10)$$

where $\Phi_{AB'}$ is the Ricci spinor, Λ is the curvature scalar spinor, $\nabla\Phi_{AB'}$ and $\nabla^2\Phi_{AB'}$ are the first and second symmetrized covariant derivatives of $\Phi_{AB'}$, $\nabla\Lambda_{AB'}$ is the first symmetrized covariant derivative of Λ and $\square\Phi_{AB'}$ and $\square\Lambda$ are the D'Alembertians of $\Phi_{AB'}$ and Λ [14].

In order to study the limits of a spacetime we first find the Cartan scalars. In general they are functions of the coordinates and the parameters of the metric. Eliminating the coordinates and parameters from as many Cartan scalars as possible (see eqs. (3.9)–(3.18) below) they can be grouped in three sets. The Cartan scalars on the first set are functions of the coordinates and of the parameters. Those on the second depend on the parameters and on the elements of the first set. Finally, the Cartan scalars on the third set are functions of the elements on the first and second ones. The expressions in the third set are parameter-free and coordinate-free, therefore they hold for any limit of the spacetime (in a given basis). The expressions in the second are coordinate-free but not parameter-free, therefore their limits depend on the limits of the Cartan scalars in the first set and the limits we choose for the parameters. The limits of the Cartan scalars on the first set can be made arbitrary with a suitable choice of coordinate system. Briefly, to obtain the limits of a spacetime in a given basis we choose a limit to the Cartan scalars in the first set and to the parameters. Then we calculate the limits of the Cartan scalars in the second and third sets. With the limit of all the Cartan scalars we have the complete characterization of the limit spacetime.

Nevertheless, this does not mean that the limit Cartan scalars one gets correspond automatically to any metric, in other words, there is no *a priori* guarantee that the Cartan scalars we obtain correspond to a Riemann geometry. It is necessary to check whether they satisfy integrability conditions [9]. This is usually very hard. Although there is no algorithm to prove that a given set of Cartan scalars correspond to a Riemann geometry, in most practical cases we can either prove they do not satisfy the integrability conditions or we can find the corresponding metric in the CLASSI database we mentioned before.

Through the steps of the Karlhede's algorithm either the dimension of the

orbit or the dimension of the isotropy group decreases, or both. Whenever the isotropy group decreases, some freedom on the choice of the basis is lost. Frame transformations could lead to new limits in the same way coordinates changes do. Therefore, fixing the frame we may loose some limits. To recover these limits we apply the lost isotropy freedom to the Cartan scalars. On the zero derivative step of the algorithm this is done with the help of the Petrov and Segre classifications. On the other steps, although a classification scheme does not exist for the derivatives, the isotropy freedom is usually of low dimension [16].

3 Nariai and O'Hanlon-Tupper spacetimes

The Nariai [17, 18, 12] and the O'Hanlon-Tupper [15] line element (with $k = 0$) and scalar field may be collectively written as

$$ds^2 = dt^2 - A^2 (dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2), \quad (3.1)$$

$$\phi = \phi_0 t^P, \quad (3.2)$$

where

$$A = A_0 t^Q, \quad (3.3)$$

$$A_0 \text{ and } \phi_0 \text{ are constants,} \quad (3.4)$$

$$Q = \begin{cases} (3\omega + 4)^{-1} \left(\omega + 1 \pm \sqrt{\frac{2\omega+3}{3}} \right) & \text{for O'Hanlon-Tupper,} \\ \frac{2+2\omega(2-\gamma)}{4+3\omega\gamma(2-\gamma)} & \text{for Nariai,} \end{cases} \quad (3.5)$$

$$P = \begin{cases} (3\omega + 4)^{-1} \left(1 \mp \sqrt{3(2\omega + 3)} \right) & \text{for O'Hanlon-Tupper,} \\ \frac{2(4-3\gamma)}{4+3\omega\gamma(2-\gamma)} & \text{for Nariai.} \end{cases} \quad (3.6)$$

Both are solutions of the Brans-Dicke field equations (2.1) with a vanishing cosmological constant. The Nariai metric is a perfect fluid solution with a equation of state given by

$$p = (\gamma - 1)\rho, \text{ where } 1 \leq \gamma \leq 2. \quad (3.7)$$

The O'Hanlon-Tupper metric is a vacuum solution. In the null tetrad

$$\begin{aligned}
\omega^0 &= 1/\sqrt{2}(dt + A dr), \\
\omega^1 &= 1/\sqrt{2}(dt - A dr), \\
\omega^2 &= 1/\sqrt{2}Ar(d\theta + i \sin(h)d\phi), \\
\omega^3 &= 1/\sqrt{2}Ar(d\theta - i \sin(h)d\phi),
\end{aligned} \tag{3.8}$$

the nonzero algebraically independent Cartan scalars (C_n) may be written as:

$$0^{\text{th}} \text{ derivative:} \quad \Phi_{00'} = 2\Phi_{11'}, \tag{3.9}$$

$$\Phi_{11'} = \frac{1}{4}Qt^{-2}, \tag{3.10}$$

$$\Phi_{22'} = 2\Phi_{11'}, \tag{3.11}$$

$$\begin{aligned}
\Lambda &= \frac{1}{4}Qt^{-2}(2Q - 1) \\
&= \Phi_{11'}(2Q - 1),
\end{aligned} \tag{3.12}$$

$$1^{\text{st}} \text{ derivative:} \quad \nabla\Phi_{00'} = \nabla\Phi_{33'}, \tag{3.13}$$

$$\nabla\Phi_{11'} = \frac{1}{3}\nabla\Phi_{33'}, \tag{3.14}$$

$$\nabla\Phi_{22'} = \frac{1}{3}\nabla\Phi_{33'}, \tag{3.15}$$

$$\begin{aligned}
\nabla\Phi_{33'} &= -\frac{1}{\sqrt{2}}Qt^{-3}(Q + 1) \\
&= -\frac{8}{\sqrt{2}}\frac{Q + 1}{\sqrt{Q}}(\phi_{11'})^{\frac{3}{2}} \\
&= -\frac{4\Phi_{11'}(\Lambda + 3\Phi_{11'})}{\sqrt{\Lambda + \Phi_{11'}}},
\end{aligned} \tag{3.16}$$

$$\nabla\Lambda_{00'} = \nabla\Lambda_{11'}, \tag{3.17}$$

$$\begin{aligned}
\nabla\Lambda_{11'} &= -\frac{1}{2\sqrt{2}}Qt^{-3}(2Q - 1) \\
&= -\frac{4}{\sqrt{2}}\frac{2Q - 1}{\sqrt{Q}}(\phi_{11'})^{\frac{3}{2}} \\
&= -\frac{4\Phi_{11'}\Lambda}{\sqrt{\Lambda + \Phi_{11'}}}.
\end{aligned} \tag{3.18}$$

We wrote as many scalars as possible as function of only few scalars. $\Phi_{11'}$ is the only scalar which is function of t and Q . Λ is function of Q and $\Phi_{11'}$ only. All the others are functions of $\Phi_{11'}$ and Λ . Therefore, once limits for $\Phi_{11'}$ and Q are chosen, the limits of the other scalars have to satisfy the coordinate-free expressions above.

Some useful information such as Petrov and Segre types and dimension of the isometry and isotropy groups can be found directly from the Cartan scalars. At the zero order derivative step it is found that the Petrov type is 0, as the Weyl spinor vanishes, and that the Segre type is $[(111),1]$. The isotropy group H_0 is the 3-dimensional group $SO(3)$. There is only one independent function of the coordinates ($\Phi_{11'}$), so $t_0 = 1$ and $d_0 = 3$. At the first derivative step we find that the isotropy group H_1 is the same as H_0 and that there is no new independent function of the coordinates. Therefore the algorithm stops.

The possible limits for $\Phi_{11'}$ are (1) 0; (2) a function of the four coordinates, $f(x^i)$; (3) a non-zero constant; or (4) ∞ . This last case is a singular limit and shall not be considered in this paper. We shall analyse the first three possibilities together with the limits of Q which are obtained from (3.5), namely¹

$$\lim_{\omega \rightarrow \infty} Q \rightarrow \begin{cases} \frac{1}{3} & \text{for O'Hanlon-Tupper,} \\ \frac{2}{3\gamma} & \gamma \neq 2 \text{ for Nariai.} \end{cases} \quad (3.19)$$

1. $\Phi_{11'} \rightarrow 0$. All Cartan scalars tend to zero, therefore the limit is Minkowski space. Regarded as a general relativity solution, this corresponds to vacuum.
2. $\Phi_{11'} \rightarrow f(x^i)$. From (3.9)–(3.18) the other Cartan scalars have finite limits. In principle this could lead to many different limit spacetimes, depending on the function $f(x^i)$. Nevertheless any such functions are related to each other by coordinate transformations, therefore the respective spacetimes are equivalent (see also [16]). These Cartan scalars correspond to the FRW metric with euclidian spatial section. The line element is given by eq. (3.1), where

$$A = A_0 t^{Q'}, \quad Q' = \text{const.} \quad (3.20)$$

¹Note that Nariai metric does not depend on ω if $\gamma = 2$. Therefore, this particular case will not be considered.

Regarded as a general relativity solution, it can be obtained from the field equations with vanishing cosmological constant and a perfect fluid matter tensor satisfying a equation of state $p = (\gamma' - 1)\rho$ where γ' and Q' are related by (see [11])

$$Q' = \frac{2}{3\gamma'}. \quad (3.21)$$

3. $\Phi_{11'} \rightarrow \text{const} \neq 0$. From the set of Cartan scalars obtained in this limit one concludes that the corresponding spacetime is conformally flat and of Segre type $[1, (111)]$, has a transitive seven-dimension group of isometry, is homogeneous and its isotropy group is the $SO(3)$. It then follows (see [11]) that the metric is a static Robertson-Walker, i.e., the Einstein metric with positive or negative tri-curvature. Nevertheless, for both these metrics, the first order Cartan scalars can be shown to be identically zero. As from eq. (3.13)–(3.18) the first order Cartan scalars of the limit are not identically zero one concludes that this set of Cartan scalars does not correspond to a riemannian geometry, i.e., it does not satisfy some integrability condition. So $\Phi_{11'} \rightarrow \text{const} \neq 0$ is not a possible limit for the metric (3.1).

This exhausts the $[(111), 1]$ Segre type limits. We shall now study the possibility of limits with a different Segre type. Similarly to the Petrov type specializations studied in [5, 16], the Segre classification satisfies a specialization scheme [20, 6, 23]. According to this scheme, the possible limits of the Segre type $[1, (111)]$ are $[(112)]$ and $[(1111)]$. The last one corresponds to $\Phi_{AB'}$ identically zero. From (3.12) this implies in $\Lambda = 0$ and therefore to Minkowski space which was already covered by our previous analysis. We proceed now with the analysis of the Segre type $[(112)]$ limits.

The standard frame we used here (see eqs. (3.9)–(3.11)) for the Segre type $[1, (111)]$ is given by the following condition:

$$\Phi_{00'} = \Phi_{22'} = 2\Phi_{11'} \quad (3.22)$$

where the other components of $\Phi_{AB'}$ vanish. The only limits that can be obtained on this frame are of the same Segre type or of Segre type $[(1111)]$. In order to study Segre type $[(112)]$ limits, the frame has to be changed. A suitable standard frame for Segre type $[(112)]$ is given by the condition [7]

$$\Phi_{22'} = 1 \quad (3.23)$$

where the other components of $\Phi_{AB'}$ vanish. To transform our previous frame to one in which the Cartan scalars might satisfy condition (3.23) on the limits we use a boost

$$\begin{pmatrix} 1/z & 0 \\ 0 & z \end{pmatrix}, \quad (3.24)$$

where z is real. The transformed Cartan scalars may be written as

$$0^{\text{th}} \text{ derivative:} \quad \tilde{\Phi}_{00'} = 2z^{-4}\Phi_{11'}, \quad (3.25)$$

$$\tilde{\Phi}_{11'} = \Phi_{11'}, \quad (3.26)$$

$$\tilde{\Phi}_{22'} = 2z^4\Phi_{11'}, \quad (3.27)$$

$$\tilde{\Lambda} = \Lambda, \quad (3.28)$$

$$1^{\text{st}} \text{ derivative:} \quad \widetilde{\nabla\Phi}_{00'} = z^{-6}\nabla\Phi_{33'}, \quad (3.29)$$

$$\widetilde{\nabla\Phi}_{11'} = 1/3z^{-2}\nabla\Phi_{33'}, \quad (3.30)$$

$$\widetilde{\nabla\Phi}_{22'} = 1/3z^2\nabla\Phi_{33'}, \quad (3.31)$$

$$\widetilde{\nabla\Phi}_{33'} = z^6\nabla\Phi_{33'}, \quad (3.32)$$

$$\widetilde{\nabla\Lambda}_{00'} = z^{-2}\nabla\Lambda_{11'}, \quad (3.33)$$

$$\widetilde{\nabla\Lambda}_{11'} = z^2\nabla\Lambda_{11'}, \quad (3.34)$$

where $\Phi_{11'}$, Λ , $\nabla\Phi_{33'}$ and $\nabla\Lambda_{11'}$ are given by (3.10), (3.12), (3.16) and (3.18).

In order to obtain the standard form (3.23), the limits of $\Phi_{11'}$ and z must be

$$\Phi_{11'} \rightarrow 0 \quad \text{and} \quad z \rightarrow \infty, \quad (3.35)$$

in such a way that the limit of the product in (3.27) is

$$2z^4\Phi_{11'} \rightarrow 1. \quad (3.36)$$

These conditions hold if and only if

$$z^4 \rightarrow \frac{1}{2\Phi_{11'}}, \quad (3.37)$$

which is a coordinate-free definition.

Using this result and the limits (3.19) in (3.28)–(3.34) we find that in the limit the nonzero Cartan scalars are

$$0^{\text{th}} \text{ derivative:} \quad \tilde{\Phi}_{22'} \rightarrow 1, \quad (3.38)$$

$$\text{1}^{\text{st}} \text{ derivative:} \quad \widetilde{\nabla} \Phi_{33'} \rightarrow -2 \frac{Q+1}{\sqrt{Q}}. \quad (3.39)$$

This set of Cartan scalars gives rise to Segre type [(112)] metrics with the isotropy groups $H_0 = H_1$ equal to a 2-parameter group of null rotation plus a 1-parameter group of spatial rotations. $d_0 = d_1 = 4$ as the Cartan scalars are constant. Therefore the spacetime is ST-homogeneous.

Searching the CLASSI database [14] for spacetimes with those Cartan scalars, we found that they are special cases of the conformally flat plane-wave class (21.38) in ref. [11], which can be written as

$$ds^2 = 2d\zeta d\bar{\zeta} - 2dudv - 2H(\zeta, \bar{\zeta}, u)du^2, \quad (3.40)$$

where H is real,

$$H_{,\zeta\zeta} = H_{,\bar{\zeta}\bar{\zeta}} = 0, \quad (3.41)$$

and where a bar and a comma denote complex conjugation and partial derivative, respectively.

In the null tetrad

$$\begin{aligned} \omega^0 &= H(H_{,\zeta\bar{\zeta}})^{-1/2}du + (H_{,\zeta\bar{\zeta}})^{-1/2}dv, \\ \omega^1 &= (H_{,\zeta\bar{\zeta}})^{1/2}du, \\ \omega^2 &= d\bar{\zeta}, \\ \omega^3 &= d\zeta \end{aligned} \quad (3.42)$$

the nonzero algebraically independent Cartan scalars (C_n) for this spacetime may be written as

$$\text{0}^{\text{th}} \text{ derivative:} \quad \tilde{\Phi}_{22'} = 1, \quad (3.43)$$

$$\text{1}^{\text{st}} \text{ derivative:} \quad \widetilde{\nabla} \Phi_{33'} = -2 \left((H_{,\zeta\bar{\zeta}})^{-1/2} \right)_{,u}, \quad (3.44)$$

$$\text{2}^{\text{nd}} \text{ derivative:} \quad \widetilde{\nabla^2} \Phi_{44'} = (H_{,\zeta\bar{\zeta}})^{-2} H_{,\zeta\bar{\zeta}uu}, \quad (3.45)$$

where we have used tilde for compatibility with the notation we have been using in the type [(112)] analysis.

With a proper choice of the function H , the limit (3.38)–(3.39) may be recovered. Indeed, equating (3.44) to the constant value in (3.39), a particular solution for H can be found, namely

$$H = 4 \left(-2 \frac{Q+1}{\sqrt{Q}} \right)^{-2} \zeta \bar{\zeta} u^{-2}. \quad (3.46)$$

All possible values of $\widetilde{\nabla}\Phi_{33'}$ in (3.39) may be obtained from this solution. Therefore, the limit Cartan scalars (3.38)–(3.39) correspond to the line element (3.40) with H given by eq. (3.46). Note that the compatibility of the second derivative (3.45) does not need to be checked, since for $\widetilde{\nabla}\Phi_{33'} = \text{const}$ the second derivative is not an independent quantity.

4 A static FRW solution as a third example

We shall now study the limits of a static FRW model with closed spatial section, which is a vacuum solution of Brans-Dicke equations with cosmological constant, found by Romero and Barros [21, 22]. Its line element and scalar field may be written as

$$ds^2 = dt^2 - A^2 \left(dr^2 + \sin^2 r (d\theta^2 + \sin^2 \theta d\varphi^2) \right), \quad (4.47)$$

$$\phi = \phi_0 e^{\pm \sqrt{\frac{2\lambda}{2\omega+3}} t}, \quad (4.48)$$

where

$$A = \sqrt{\frac{2\omega+3}{\lambda(\omega+1)}}, \quad (4.49)$$

$$\phi_0 \text{ is a constant}, \quad (4.50)$$

and λ is the cosmological constant. In the null tetrad

$$\begin{aligned} \omega^0 &= 1/\sqrt{2}(dt + A dr), \\ \omega^1 &= 1/\sqrt{2}(dt - A dr), \\ \omega^2 &= 1/\sqrt{2}A \sin(r)(d\theta + i \sin(h)d\varphi), \\ \omega^3 &= 1/\sqrt{2}A \sin(r)(d\theta - i \sin(h)d\varphi), \end{aligned} \quad (4.51)$$

the nonzero algebraically independent Cartan scalars may be written as

$$0^{\text{th}} \text{ derivative:} \quad \Phi_{00'} = 2\Phi_{11'}, \quad (4.52)$$

$$\Phi_{11'} = \frac{1}{4}A^{-2}, \quad (4.53)$$

$$\Phi_{22'} = 2\Phi_{11'}, \quad (4.54)$$

$$\Lambda = \Phi_{11'}. \quad (4.55)$$

This is a Petrov type 0 and Segré type $[1,(111)]$ metric. As there is no function of the coordinates among the Cartan scalars, its a spacetime homogeneous metric. Its isometry group is seven-dimensional and the isotropy subgroup is the $SO(3)$. The first order Cartan scalars vanish identically, therefore the Karlhede algorithm stops. From (4.49), as $\omega \rightarrow \infty$,

$$A \rightarrow \sqrt{\frac{2}{\lambda}}. \quad (4.56)$$

The limit of the Cartan scalars are given by substituting A in (4.52)–(4.55) for (4.56). This is nothing but the Einstein static model of the universe, which is a solution of the general relativity field equations with cosmological constant (equals to half of ours) and a matter distribution corresponding to dust.

In this case no other limit is possible since the Cartan scalars do not depend on the coordinates.

5 Conclusion

It was found that the Nariai metric may have three possible limits as $\omega \rightarrow \infty$, namely: the Minkowski space, the FRW metric with euclidian spatial section and the conformally flat plane-wave solution (3.46). Regarded as solutions of the general relativity equations, these correspond to three different matter distributions: vacuum, perfect fluid with a equation of state $p = (\gamma' - 1)\rho$ and a Segre type $[(112)]$ configuration. The matter distribution corresponding to the first and the third metrics differ from that of Nariai's solution of Brans-Dicke theory. On the other hand, the second matter distribution is the same perfect fluid as in Nariai as can be seen from eqs. (3.19) and (3.21).

The O’Hanlon-Tupper metric has the same limits as Nariai’s. Nevertheless, it is a vacuum solution of the Brans-Dicke field equations. Therefore, the first limit, namely, the Minkowski space, has the same matter configuration (vacuum), while this does not occur with the other two limits.

The situation with the static vacuum solution of the Brans-Dicke equations with cosmological constant discussed in section 4 is quite different. It has just one limit, namely, the Einstein static metric, which regarded as a general relativity solution cannot be obtained from the vacuum Einstein’s equation, since the Ricci spinor ($\Phi_{AB'}$) does not vanish.

In other words, it seems that, in general, Brans-Dicke solutions do not have a unique limit as $\omega \rightarrow \infty$. Thus, to say that a given Brans-Dicke spacetime tends to a spacetime of general relativity may sound meaningless. Moreover, it may even happen, as the example in section 4 explicitly shows, that none of the limits corresponds to a general relativity solution with the same matter distribution.

Acknowledgments

The authors thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for financial support. F. M. Paiva also thanks Marcelo Rebouças, Antonio Teixeira and Graham Hall for many useful discussions and the Departamento de Física da Universidade Federal da Paraíba for the kind hospitality during the final stage of this work.

References

- [1] Åman, J. E. (1987), “*Manual for CLASSI - Classification Programs for Geometries in General Relativity (Third Provisional Edition)*”, University of Stockholm Report.
- [2] Brans, C. and Dicke, R. H. (1961), “*Mach’s Principle and a Relativistic Theory of Gravitation*”, Phys. Rev. **124**, 925–935.
- [3] Cartan, E. (1951), “*Leçons sur la Géométrie des Espaces de Riemann*”, Gauthier-Villars, Paris.

- [4] Frick, I. (1977), “*SHEEP Users Guide*”, Institute of Theoretical Physics, University of Stockholm Report 77–14.
- [5] Geroch, R. (1969), “*Limits of Spacetimes*”, Commun. Math. Phys. **13**, 180–193.
- [6] Hall, G. S. (1985), “*The Energy-Momentum Tensor in General Relativity*”, Lecture notes, Belatonszéplak, Hungary.
- [7] Joly, G. C. and MacCallum, M. A. H. (1990), “*Computer-aided classification of the Ricci tensor in general relativity*”, Class. Quantum Grav. **7**, 541–556.
- [8] Karlhede, A. (1980), “*A Review of the Geometrical Equivalence of Metrics in General Relativity*”, Gen. Rel. Grav. **12**, 693–707.
- [9] Karlhede, A. and Lindström, U. (1983), “*Finding Space-Time Geometries without Using a Metric*”, Gen. Rel. Grav. **15**, 597–610.
- [10] Koutras, A. (1992), “*A spacetime for which the Karlhede invariant classification requires the fourth covariant derivative of the Riemann tensor*”, Class. Quantum Grav. **9**, L143–L145.
- [11] Kramer, D., Stephani, H., MacCallum, M. and Herlt, E. (1980), “*Exact Solutions of Einstein’s Field Equations*”, Cambridge U. P., Cambridge.
- [12] Lorenz, D. (1983), “*Exact Brans-Dicke-Bianchi solutions*”, in Lectures Notes in Physics 205 — Solutions of Einstein’s Equations: Techniques and Results, Springer-Verlag.
- [13] MacCallum, M. A. H. and Åman, J. E. (1986), “*Algebraically independent n^{th} derivatives of the Riemann curvature in general relativity*”, Class. Quantum Grav. **3**, 1133–1141.
- [14] MacCallum, M. A. H. and Skea, J. E. F. (1993), “*SHEEP: A Computer Algebra System for General Relativity*”, in “Algebraic Computing in General Relativity: Lecture Notes from the First Brazilian School on Computer Algebra”, Vol. 2, edited by M. J. Rebouças and W. L. Roque, Oxford U. P., Oxford.

- [15] O’Hanlon, J. and Tupper, B. O. J. (1972), “*Vacuum-Field Solutions in the Brans-Dicke Theory*”, *Nuovo Cimento* **B7**, 305–312.
- [16] Paiva, F. M., Rebouças, M. J. and MacCallum, M. A. H. (1993), “On limits of spacetimes – a coordinate-free approach”, *Class. Quantum Grav.* To appear.
- [17] Nariai, H. (1968), “*On the Green’s Functions in an Expanding Universe and its Role in the Problem of Mach’s Principle*”, *Prog. Theor. Phys.* **40**, 49–59.
- [18] Nariai, H. (1969), “*Gravitational Instability in Brans-Dicke Cosmology*”, *Prog. Theor. Phys.* **42**, 544–554.
- [19] Penrose, R. (1960), “*A Spinor Approach to General Relativity*”, *Ann. Phys. (NY)* **10**, 171–201.
- [20] Penrose, R. (1972), “*Spinor classification of energy tensors*”, in *Gravitation: Problems, Prospects (Dedicated to the memory of A. Z. Petrov)*. Izdat, Naukova Dumka, Kiev.
- [21] Romero, C. and Barros, A. (1993), “Does Brans-Dicke theory of gravity go over to general relativity when $\omega \rightarrow \infty$?”, *Phys. Lett.. A* **173**, 243.
- [22] Romero, C. and Barros, A. (1993), “*Brans-Dicke vacuum solutions and the cosmological constant: a qualitative analysis*”, *Gen. Rel. Grav.* To appear.
- [23] Sánchez, A. R. G., Plebański, J. F. and Przanowski (1991), “*Deformations of algebraic types of the energy-momentum tensor*”, *J. Math. Phys.* **32**, 2838–2847.
- [24] Weinberg, S. (1972), “*Gravitation and Cosmology*”, New York, John Wiley and Sons.